

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Understanding B Lymphocyte Development: A Long Way to Go

---

Malavika Bhattacharya

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.79663>

---

## Abstract

B lymphocytes play a significant role in both antigen-dependent and antigen-independent pathways. Although significant progress has been made in context of delineating the pathways that lead to their development, maturity and differentiation, their detailed mechanisms are yet to be dissected out completely. This chapter is aimed towards summarising the knowledge that has been gained till date and identifying the areas that need to be addressed in future research work. Overall, the chapter is planned in a sequential way to guide the reader through the processes of B cell development and the various latest findings that have improved our understanding of this vital physiological system.

**Keywords:** B lymphocyte, bone marrow, marginal zone, antigen, spleen, germinal centre

---

## 1. Introduction

B lymphocytes play vital role in maintaining the normal immunologic functions of the body. Their functions range from producing antibodies to presenting antigens. They are also involved with productions of several regulatory cytokines, such as IL-2, IL-4, IL-6, IL-10, IL-12, TGF- $\beta$ 1, TNF and IFN- $\gamma$ . Each of these functions are further fine tuned by the fact that they are dependent upon several factors, including the B lymphocyte subsets, their location and the type of stimuli that is encountered by the specific B cell subset in that particular environment. Understanding all these components of B cell development is not only required for getting a better picture of B cell biology, but is also necessary for understanding various immunologic anomalies that lead to disorders. They are also important towards generating effective B cell based therapies. This chapter begins with explanation of the overall pathway

---

of B cell development and the organs that are involved in its various stages. This is followed by discussion on the role of gene rearrangements in the entire process. Subsequently, the role of various transcription factors has been addressed.

## 2. Steps in B lymphocyte developmental pathway

### 2.1. Bone marrow-dependent stages

During foetal life, B cells are generated in the foetal liver. Subsequently in the adults, they are produced by differentiation of haematopoietic cells (HSCs) in the bone marrow [1]. Most of the stages of B lymphocyte development take place in this primary lymphoid organ. The pluripotent HSCs gradually differentiate into progenitors, which have increasingly lower potency. Initially, they form a population of cells that are known as multipotent progenitors (MPPs). These progenitors, in turn, give rise to two main progenitor populations: common granulocyte/megakaryocyte/granulocyte progenitor (CFU-GEMM) and early lymphoid progenitor (ELP). CFU-GEMMs subsequently develop into cells that have either myeloid or erythroid potential. On the other hand, cells with lymphoid potential arise from ELPs. Thus, CFU-GEMMs are the primary source of those elements of blood that are non-lymphoid in nature, whereas the lymphoid elements originate from ELPs. Two major precursors arise from the ELPs, common lymphocyte progenitor (CLP) and early T-lineage precursor (ETP). Both Pre-NK cells and Pre-B cells develop from CLPs, which eventually give rise to NK cells and B cells, respectively. T cells derive from thymocytes, which are generated by differentiation of the ETPs. The CLPs give rise to early Pro-B cells first. They mature to form the late Pro-B cells, which eventually develop into Pre-B cells. Immature B cells arise from these Pre-B cells and they leave the bone marrow to enter into the secondary lymphoid organs. Subsequent stages of B cell development primarily continue in the spleen. During this entire period of maturation, the various B cell subpopulations are found to migrate within the bone marrow, keeping in touch with the stromal cells. Initially, the progenitor cells having highest potency lie in the endosteum. This region is located near the inner surface of the bone. As the B cell progenitors mature to give rise to cell types that have less potency and are more committed towards the B cell fate, they start migrating towards the central sinus of the bone marrow cavity. During this entire process, these maturing cell populations are reported to remain in contact with the reticular stromal cells, which are believed to provide indispensable signals for migration and maturation. This process continues till the developing cell reaches the stage of Immature B cell [2–4].

### 2.2. Role of bone marrow cell populations in B cell development

In addition to HSCs, B lymphocytes (in various stages of development) and plasma cells (PCs), the bone marrow consists of specialised cells that have multiple roles in various stages of this developmental process. Together they form the “niches” that are vital for normal functioning of the different systems associated with the bone marrow. The most important components of these “niches” are stromal cells and regulatory T cells. Out of these, the stromal cells form an

extensive network of non-lymphoid connective tissue in the bone marrow [5, 6]. They serve dual functions of forming adhesive contacts with developing lymphocytes and providing cytokines/chemokines/growth factors to them as per requirement. In context of maintenance of plasma cells, earlier reports have indicated that they are provided with survival signals such as CXCL13, IL-6, APRIL and BL $\gamma$ S by the stromal cells [7]. In addition, it has been found recently that the regulatory T cells present in the bone marrow play vital role in maintaining plasma cell pool [8]. The dendritic cells (DCs), present as perivascular clusters in the bone marrow, have also been reported to provide signals that are vital for B lymphocytes [9]. Megakaryocytes [10], eosinophils [11] as well as basophils [12] resident in the bone marrow have also been found to play role in maintaining plasma cell pools.

### 2.3. Spleen-dependent stages

The immature B cell undergoes final stages of development in the spleen to form mature B cells. The spleen primarily consists of red pulp, white pulp and marginal zone. The red pulp is made up of large, blood-filled sinuses and serves as the blood-filtering system of the spleen. In addition, the splenic macrophages play important role in recycling of iron. The white pulp is organised in line with the lymph nodes and consists of lymphoid sheaths having distinct B-cell and T-cell compartments. The marginal zone is a layer of highly specialised cells that surrounds the white pulp [13]. It plays a very important role in immunity because those haematopoietic cells that remain in circulation (as part of the surveillance mechanism) need to be able to migrate through blood and lymphatic systems continuously. The marginal zone plays a vital role in this process due to its strategic location in the spleen. It has been observed that G-protein linked receptors are involved in the signalling process that is responsible for active transport of B- and T-lymphocytes to-and-from the white pulp [14]. The specialised cells that constitute the marginal zone include two subsets of macrophages, the marginal-zone macrophages and the marginal-zone metallophilic macrophages. The first subset is present as an outer ring and express SIGNR1 (a C-type lectin) [15–17] and MARCO (a type I scavenger receptor) [18]. The second subset is present as an inner ring, lies closer to the white pulp and expresses SIGLEC-I (an adhesion molecule) [19]. A specialised B-cell population, known as marginal zone B cell, and DCs are located in between these two rings of macrophages [20, 21]. **Figure 1** shows the major cell populations that are generated in the bone marrow and peripheral lymphoid organs during the process of B cell development.

It has also been reported that the antigens are encountered by the mature B cells in the lymphoid follicles. This process is aided by T cells present in the germinal centres. All the subsequent stages of B cell development, including generation of various Ig isotypes, class switching and somatic hypermutation, contribute towards diversification of the antibody repertoire [22].

On activation by cognate antigen, the activated B cell can either differentiate into antibody-secreting plasma cells/plasma blasts or get recruited into a specialised region known as the germinal centre (GC). Those activated B cells that enter the GC subsequently undergo several rounds of proliferation, class-switching and affinity maturation. Thereafter, the GC B cells that have completed these steps successfully give rise to either long-lived plasma cells or




Figure 1. Major stages of B cell development.




Figure 2. Major stages after activation of B cells.

memory B cells [23]. When the B cells undergo these changes in the GC, their transcriptional repertoire also undergoes huge transformation [24]. **Figure 2** shows the various stages that are developed after activation of B cells.



### 3. B cell subsets

In general, three subsets of B cells derive from naive B cells, B-1 B cells, follicular B cells and marginal zone (MZ) B cells. Furthermore, B-1B cells form two subsets, B-1a and B-1 b B cells [25]. All the subsets can be clearly identified on the basis of their surface markers. These surface markers can also be used to identify their progenitor populations. **Table 1** shows the various stages of B cell development and the most prominent cell surface markers that are used to identify them. Each one of these subsets maintains distinct location(s) and function(s). All these subsets of B cells produce functionally important antibodies. However, they vary in huge terms in reference to their origin and function [26–28].

#### 3.1. B-1 B cells

B-1 B cells have not been successfully studied in mammals, including humans [29]. Thus, most of the findings are based on studies performed in mice. The most interesting finding from these studies is that although the progenitors of B-1 a and B-1 b cells are distinct, they are found to occupy the same locations, namely the pleural and peritoneal cavities. In addition, it has been observed that the environment offered by these cavities plays a significant role in

Developmental stage	Mouse cell surface marker		Human cell surface marker	
	Positive	Negative	Positive	Negative
Pre-Pro B	CD43, B220, Flt3, CD93, CXCR4, IL7R	CD19, cKit <sup>low</sup>	CD34, CD38, CD10	cKit <sup>low</sup>
Pro B	CD19, CD43, B220, IL7R	Flt3, ckit <sup>low</sup>	CD19, CD34, CD38, CD10, CD24, IL7/3R	ckit <sup>low</sup>
Pre B	CD19, B220, IL7R, CD24, Siglec-G	CD43 <sup>low</sup> , ckit	CD19, CD10, CD20, CD24, CD38, IL3/4/7R	CD34, ckit
Immature B	CD19, CD43, B220, CD24	CD43, CD23	CD19, CD10, CD20, CD21, CD40, IL4R, CD24 <sup>high</sup> , CD38 <sup>high</sup>	ckit, CD27, IL7R
Transitional B	B220, CD19, CD24, CD93, CD21 <sup>var</sup> , CD23 <sup>var</sup>		CD5, CD21, CD19, CD20, CD24 <sup>high</sup> , CD38 <sup>high</sup>	CD27, CD10 <sup>low</sup>
Marginal Zone B	B220, CD9, CD1d, CD21 <sup>high</sup> , CD35 <sup>high</sup>	CD93, CD23	CD19, CD20, CD1c, CD21 <sup>high</sup> , CD27 <sup>var</sup>	
Regulatory B	CD19, CD5, CD24, TIM, CD1d <sup>high</sup>	CD62L, CD93 <sup>var</sup>	CD19, CD5, CD21, CD1d <sup>high</sup> , CD24 <sup>high</sup>	CD27 <sup>var</sup>
Follicular B	B220, CD19, CD23, CD38, CD22	CD93, CD1d <sup>low</sup> , CD21/35 <sup>low</sup>	CD19, CD20, CD21, CD24, CD23, CD22	CD27, CD10, CD24 <sup>low</sup> , CD38 <sup>low</sup>
Activated B	B220, Flt3, CD27, CD80, MHC II <sup>high</sup>	CXCR4, CD138	CD19, CD20, CD27, CD25, CD69, CD30, CD135	
Germinal Center B	CD19, CD20, Siglec2, GL7, CD37	CD93, CD38 <sup>low</sup>	CD19, CD10, CD23, CD27, CD20, CD38 <sup>high</sup>	CD24 <sup>low</sup>
Plasmablast	CD138, CD19, MHC II, CXCR4	Flt3, B220 <sup>low</sup>	CD19, MHC II, CD269, CD38 <sup>high</sup> , CD27 <sup>high</sup>	CD138, CD20
Plasma cell (short lived)	CD138, CD93, CXCR4 <sup>high</sup>	CD19, B220 <sup>low</sup> , CD38 <sup>low</sup> , MHC II <sup>low</sup>	CD138, CD269, CD27, CXCR4, CD38 <sup>high</sup>	CD20, CD19 <sup>low</sup> , MHC II <sup>low</sup>
Plasma cell (long lived)	CD138, CXCR4 <sup>high</sup>	CD19, B220 <sup>low</sup> , CD38 <sup>low</sup> , MHC II	CD138, CD269, CD27, CXCR4, CD38 <sup>high</sup>	CD20, CD19 <sup>low</sup> , MHC II
Memory B	B220, CD38 <sup>var</sup> , CD80 <sup>var</sup> , CD62L <sup>var</sup> , CD95 <sup>low</sup>		CD19, CD40, CD20, CXCR4/5/7, CD27 <sup>var</sup>	CD38, CD23 <sup>low</sup>

**Table 1.** Most prominent cell surface markers used to identify various stages of B cell development.

shaping the functional characteristics of these B cell subsets. The milieu of these cavities also influences the functional characteristics of B-2 cells that reside there, although in low numbers [30, 31].

### 3.2. Follicular B cells

Several studies have shown that the naive, mature peripheral follicular B cells reside in two main niches during their circulation/recirculation through the bone marrow. Out of these, the “follicular niche” present in the spleen/lymph nodes/Peyer’s patches is the main site that is occupied by these cells. These “follicular sites” are thought to play important role in those immune responses against protein antigens, which are T cell-dependent [25].

In addition, some follicular B cells have also been reported to home in the bone marrow [32]. The site of their homing has been termed as “perisinusoidal niche” and consists of a part of population of the same circulating follicular B cells that are found in the secondary lymphoid organs mentioned earlier. Interestingly, the follicular B cells residing in the bone marrow are involved in T cell-independent immune responses against microbial pathogens harboured by blood [33].

### 3.3. Marginal zone (MZ) B cells

MZ B cells mainly home near the marginal sinus of the spleen. This process is facilitated by the molecules SIP 1 and SIP 3, which are receptors for sphingosine-1-phosphate [34–36]. These cells are mainly involved in T cell-independent immune responses against blood-borne microbes [34]. It has also been reported that these MZ B cells can transport pathogens from the marginal sinus to the splenic follicles, sites where the follicular B cells reside [21, 37]. Moreover, a few *in vitro* studies have shown that these MZ B cells might also be contributing towards T cell-dependent pathways of antigen recognition and subsequent immune responses. Some reports have demonstrated that they may have better capability than follicular B cells in context of activation of T cells [38, 39].

## 4. Contributions of various players towards B cell development and selection

### 4.1. Pathways involved in B cell maturation stages in bone marrow

As mentioned earlier, B cell development starts from haematopoietic stem cells (HSCs) in the bone marrow and continues either at the same site or in the peripheral sites. Two cellular pathways are believed to be involved in formation of mature B cells from T2-like cells, one in the bone marrow, and the other in the peripheral sites [25]. Thus, the population of mature B cells present in the bone marrow is heterogeneous in nature. It has been observed that one population of these cells is characterised by sIgM<sup>high</sup> IgD<sup>low</sup> CD23<sup>-</sup>, and do not respond either to BLγS/BAFF or multivalent antigens. In contrast, another population of these cells is sIgM<sup>high</sup>

IgD<sup>high</sup> CD23<sup>+</sup>, and responds to both BL $\gamma$ S/BAFF and multivalent antigens [32, 40]. The exact mechanisms that are responsible for this differential developmental pathways and their significance is yet to be understood completely.

#### **4.2. Pathways involved in B cell maturation at the peripheral tissues**

Once immature B cells form, they generally migrate to the peripheral sites. These immature B cells have very short half-lives and on engagement with BCRs, they tend to undergo apoptosis instead of proliferation [41, 42]. The immature B cells are also referred to as “transitional B cells” and can be further subdivided into three subsets; T1, T2 and T3 [41, 43, 44]. CD93/AA4 (the B-lineage precursor marker) is expressed in all of these “transitional B cell” subsets. However, they display differential expression of IgM and CD23 on their surfaces that is exploited to identify them. T1 cells are characterised by IgM<sup>high</sup> CD23<sup>-</sup>, T2 cells by IgM<sup>high</sup> CD23<sup>+</sup> and T3 cells by IgM<sup>low</sup> CD23<sup>-</sup> [41]. All data till date suggest that T2-like cells give rise to mature B cells, either in the bone marrow or in the peripheral sites.

#### **4.3. Tolerance and B cell development in peripheral sites**

Tolerance to self-reactive antigens can take place by any of the existing three mechanisms: deletion, editing or anergy. In spite of significant number of studies addressing this aspect, it is not yet clear whether this tolerance is achieved by negative selection of self-reactive B cell clones or failure of positive selection. Several studies have suggested that local levels of BAFF may influence this decision [45, 46].

In humans, majority (around 55–75%) of immature B cells have been found to be self-reactive [47], indicating that clonal deletion may serve as one of the main mechanisms of elimination of these self-reactive cell populations. However, no such experimental data is available from mice. In addition, receptor editing has also been found to contribute towards elimination of such self-reactive immature B cells. The process of anergy, although a bit controversial, has emerged as the third mechanism involved in removal of self-reactive B cells [48].

#### **4.4. Pathways involved in B cell maturation stages in the spleen**

Spleen is the main site for positive selection of non-self reactive B cell clones. It has been reported that survival of the peripheral B cell populations is dependent upon continuous signalling through B Cell Receptor (BCR). It holds true for both populations, follicular B cells as well as marginal zone (MZ) B cells [49, 50]. Although a wide variety of V-gene segments are expressed by both follicular and MZ B cell populations, it has been observed that in IgH transgenic mice, those immature B cells that are specific for phosphorylcholine give rise to MZ B cells [51]. In another study, it has been observed that deletion of RAG2 in adult mice results in selective retention of MZ B cells over follicular B cells [52].

In addition to BCR, Notch2 signalling plays a significant role in MZ B cell development [53, 54]. Notch2 is a member of Notch family of receptors and ligands [55]. Interaction between Notch2 and DL1 has been found to be responsible for developing a unique MZB cell niche [54]. Moreover,









## Notes

My sincere apologies go to all those researchers who are doing pioneering work in this field too, but whom I have not been able to quote in my article due to space constraint.

## Author details

Malavika Bhattacharya

Address all correspondence to: malavikab@gmail.com

Department of Biotechnology, Techno India University, Kolkata, West Bengal, India

## References

- [1] Dorshkind K. Multilineage development from adult bone marrow cells. *Nature Immunology*. 2002;**3**:311-313
- [2] Mansson R, Hultquist A, Luc S, et al. Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. *Immunity*. 2007;**26**:407-419
- [3] Laiosa CV, Stadtfeld M, Graf T. Determinants of lymphoid-myeloid lineage diversification. *Annual Review of Immunology*. 2006;**24**:705-738
- [4] Nagasawa T. Microenvironmental niches in the bone marrow required for B-cell development. *Nature Reviews. Immunology*. 2006;**6**:107-116
- [5] Tokoyoda K, Egawa T, Sugiyama T, et al. Cellular niches controlling B lymphocyte behavior within bone marrow during development. *Immunity*. 2004;**20**:707-718
- [6] Zehentmeier S, Roth K, Cseresnyes Z, et al. Static and dynamic components synergize to form a stable survival niche for bone marrow plasma cells. *European Journal of Immunology*. 2014;**44**:2306-2317
- [7] Roth K, Oehme L, Zehentmeier S, et al. Tracking plasma cell differentiation and survival. *Cytometry. Part A*. 2014;**85**:15-24
- [8] Zaretsky AG, Konradt C, De' pis F, et al. T regulatory cells support plasma cell populations in the bone marrow. *Cell Reports*. 2017;**18**:1906-1916
- [9] Sapozhnikov A, Pewzner-Jung Y, Kalchenko V, et al. Perivascular clusters of dendritic cells provide critical survival signals to B cells in bone marrow niches. *Nature Immunology*. 2008;**9**:388-395
- [10] Winter O, Moser K, Mohr E, et al. Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. *Blood*. 2010;**116**:1867-1875

- [11] Chu VT, Fröhlich A, Steinhäuser G, et al. Eosinophils are required for the maintenance of plasma cells in the bone marrow. *Nature Immunology*. 2011;**12**:151-159
- [12] Rodriguez Gomez M, Talke Y, Goebel N, et al. Basophils support the survival of plasma cells in mice. *Journal of Immunology*. 2010;**185**:7180-7185
- [13] Mebius RE, Kraal G. Structure and function of the spleen. *Nature Reviews. Immunology*. 2005;**5**:606-616
- [14] Cyster JG, Goodnow CC. Pertussis toxin inhibits migration of B and T lymphocytes into splenic white pulp cords. *The Journal of Experimental Medicine*. 1995;**182**:581-586
- [15] Kang YS et al. The C-type lectin SIGN-R1 mediates uptake of the capsular polysaccharide of *Streptococcus pneumoniae* in the marginal zone of mouse spleen. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;**101**:215-220
- [16] Kang YS, Yamazaki S, Iyoda T, et al. SIGN-R1, a novel C-type lectin expressed by marginal zone macrophages in spleen, mediates uptake of the polysaccharide dextran. *International Immunology*. 2003;**15**:177-186
- [17] Geijtenbeek TB, Groot PC, Nolte MA, et al. Marginal zone macrophages express a murine homologue of DC-SIGN that captures blood-borne antigens in vivo. *Blood*. 2002;**100**:2908-2916
- [18] Elomaa O et al. Cloning of a novel bacteria-binding receptor structurally related to scavenger receptors and expressed in a subset of macrophages. *Cell*. 1995;**80**:603-609
- [19] Munday J, Floyd H, Crocker PR. Sialic acid binding receptors (siglecs) expressed by macrophages. *Journal of Leukocyte Biology*. 1999;**66**:705-711
- [20] Yu P et al. B cells control the migration of a subset of dendritic cells into B cell follicles via CXC chemokine ligand 13 in a lymphotoxin-dependent fashion. *Journal of Immunology*. 2002;**168**:5117-5123
- [21] Martin F, Kearney JF. Marginal-zone B cells. *Nature Reviews. Immunology*. 2002;**2**:323-335
- [22] LeBien TW, Tedder TF. B lymphocytes: How they develop and function. *Blood*. 2008;**112**:1570-1580
- [23] Nutt SL, Hodgkin PD, Tarlinton DM, Corcoran LM. The generation of antibody-secreting plasma cells. *Nature Reviews. Immunology*. 2015;**15**:160-171
- [24] Shi W, Liao Y, Willis SN, et al. Transcriptional profiling of mouse B cell terminal differentiation defines a signature for antibody-secreting plasma cells. *Nature Immunology*. 2015;**16**:663-673
- [25] Allman D, Pillai S. Peripheral B cell subsets. *Current Opinion in Immunology*. 2008;**20**:149-157
- [26] Kantor AB, Herzenberg LA. Origin of murine B cell lineages. *Annual Review of Immunology*. 1993;**11**:501-538



- [27] Hardy RR, Hayakawa K. B cell development pathways. *Annual Review of Immunology*. 2001;**19**:595-621
- [28] Baumgarth N. The double life of a B-1 cell: Self-reactivity selects for protective effector functions. *Nature Reviews. Immunology*. 2011;**11**:34-46
- [29] Thiriot A, Drapier AM, Vieira P, et al. The Bw cells, a novel B cell population conserved in the whole genus *Mus*. *Journal of Immunology*. 2007;**179**:6568-6578
- [30] Hastings WD, Tumang JR, Behrens TW, Rothstein TL. Peritoneal B-2 cells comprise a distinct B-2 cell population with B-1b-like characteristics. *European Journal of Immunology*. 2006;**36**:1114-1123
- [31] Berberich S, Forster R, Pabst O. The peritoneal microenvironment commits B cells to home to body cavities and the small intestine. *Blood*. 2007;**109**:4627-4634
- [32] Cariappa A, Chase C, Liu H, et al. Naive recirculating B cells mature simultaneously in the spleen and bone marrow. *Blood*. 2007;**109**:2339-2345
- [33] Cariappa A, Mazo IB, Chase C, et al. Perisinusoidal B cells in the bone marrow participate in T-independent responses to blood-borne microbes. *Immunity*. 2005;**23**:397-407
- [34] Cinamon G, Matloubian M, Lesneski MJ, et al. Sphingosine 1-phosphate receptor 1 promotes B cell localization in the splenic marginal zone. *Nature Immunology*. 2004;**5**:713-720
- [35] Vora KA, Nichols E, Porter G, et al. Sphingosine 1-phosphate receptor agonist FTY720-phosphate causes marginal zone B cell displacement. *Journal of Leukocyte Biology*. 2005;**78**:471-480
- [36] Cinamon G, Zachariah MA, Lam OM, et al. Follicular shuttling of marginal zone B cells facilitates antigen transport. *Nature Immunology*. 2008;**9**:54-62
- [37] Ferguson AR, Youd ME, Corley RB. Marginal zone B cells transport and deposit IgM-containing immune complexes onto follicular dendritic cells. *International Immunology*. 2004;**16**:1411-1422
- [38] Attanavanich K, Kearney JF. Marginal zone, but not follicular B cells, are potent activators of naive CD4 T cells. *Journal of Immunology*. 2004;**172**:803-811
- [39] Song H, Cerny J. Functional heterogeneity of marginal zone B cells revealed by their ability to generate both early antibody-forming cells and germinal centers with hypermutation and memory in response to a T-dependent antigen. *The Journal of Experimental Medicine*. 2003;**198**:1923-1935
- [40] Lindsley RC, Thomas M, Srivastava B, Allman D. Generation of peripheral B cells occurs via two spatially and temporally distinct pathways. *Blood*. 2007;**109**:2521-2528
- [41] Allman D, Lindsley RC, DeMuth W, et al. Resolution of three nonproliferative immature splenic B cell subsets reveals multiple selection points during peripheral B cell maturation. *Journal of Immunology*. 2001;**167**:6834-6840

- [42] Norvell A, Mandik L, Monroe JG. Engagement of the antigen-receptor on immature murine B lymphocytes results in death by apoptosis. *Journal of Immunology*. 1995; **154**:4404-4413
- [43] Loder F, Mutschler B, Ray RJ, et al. B cell development in the spleen takes place in discrete steps and is determined by the quality of B cell receptor-derived signals. *The Journal of Experimental Medicine*. 1999; **190**:75-89
- [44] Srivastava B, Quinn WJ 3rd, Hazard K, et al. Characterization of marginal zone B cell precursors. *The Journal of Experimental Medicine*. 2005; **202**:1225-1234
- [45] Lesley R, Xu Y, Kalled SL, et al. Reduced competitiveness of autoantigen-engaged B cells due to increased dependence on BAFF. *Immunity*. 2004; **20**:441-453
- [46] Thien M, Phan TG, Gardam S, et al. Excess BAFF rescues self-reactive B cells from peripheral deletion and allows them to enter forbidden follicular and marginal zone niches. *Immunity*. 2004; **20**:785-798
- [47] Wardemann H, Yurasov S, Schaefer A, et al. Predominant autoantibody production by early human B cell precursors. *Science*. 2003; **301**:1374-1377
- [48] Cyster JG, Hartley SB, Goodnow CC. Competition for follicular niches excludes self-reactive cells from the recirculating B-cell repertoire. *Nature*. 1994; **371**:389-395
- [49] Kraus M, Alimzhanov MB, Rajewsky N, Rajewsky K. Survival of resting mature B lymphocytes depends on BCR signaling via the Igalpha/beta heterodimer. *Cell*. 2004; **117**:787-800
- [50] Cariappa A, Tang M, Parng C, et al. The follicular versus marginal zone B lymphocyte cell fate decision is regulated by Aiolos, Btk, and CD21. *Immunity*. 2001; **14**:603-615
- [51] Martin F, Kearney JF. Positive selection from newly formed to marginal zone B cells depends on the rate of clonal production, CD19, and btk. *Immunity*. 2000; **12**:39-49
- [52] Hao Z, Rajewsky K. Homeostasis of peripheral B cells in the absence of B cell influx from the bone marrow. *The Journal of Experimental Medicine*. 2001; **194**:1151-1164
- [53] Saito T, Chiba S, Ichikawa M, et al. Notch2 is preferentially expressed in mature B cells and indispensable for marginal zone B lineage development. *Immunity*. 2003; **18**:675-685
- [54] Hozumi K, Negishi N, Suzuki D, et al. Delta-like 1 is necessary for the generation of marginal zone B cells but not T cells in vivo. *Nature Immunology*. 2004; **5**:638-644
- [55] Maillard I, Fang T, Pear WS. Regulation of lymphoid development, differentiation, and function by the Notch pathway. *Annual Review of Immunology*. 2005; **23**:945-974
- [56] Moran ST, Cariappa A, Liu H, et al. Synergism between NF-kappa B1/p50 and Notch2 during the development of marginal zone B lymphocytes. *Journal of Immunology*. 2007; **179**:195-200
- [57] Sasaki Y, Derudder E, Hobeika E, et al. Canonical NF-kappaB activity, dispensable for B cell development, replaces BAFF-receptor signals and promotes B cell proliferation upon activation. *Immunity*. 2006; **24**:729-739

- [58] Thomas M, Calamito M, Srivastava B, et al. Notch activity synergizes with B-cell-receptor and CD40 signaling to enhance B-cell activation. *Blood*. 2007;**109**:3342-3350
- [59] Hozumi N, Tonegawa S. Evidence for somatic rearrangement of immunoglobulin genes coding for variable and constant regions. *Proceedings of the National Academy of Sciences of the United States of America*. 1976;**73**(10):3628-3632
- [60] Tonegawa S. Somatic generation of antibody diversity. *Nature*. 1983;**302**(5909):575-581
- [61] Alt FW, Blackwell TK, Yancopoulos GD. Development of the primary antibody repertoire. *Science*. 1987;**238**:1079-1087
- [62] Rolink A, Melchers F. Molecular and cellular origins of B lymphocyte diversity. *Cell*. 1991;**66**:1081-1094
- [63] Lansford R, Okada A, Chen J, et al. Mechanism and control of immunoglobulin gene rearrangement. In: Hames BD, Glover DM, editors. *Molecular Immunology*, 1996; *Frontiers in Molecular Biology*. 2nd ed. Oxford: Oxford University Press. pp. 1-100
- [64] Spanopoulou E, Roman CAJ, Corcoran LM, et al. Functional immunoglobulin transgenes guide ordered B-cell differentiation in Rag-1-deficient mice. *Genes & Development*. 1994;**8**:1030-1042
- [65] Young F, Ardman B, Shinkai Y, et al. Influence of immunoglobulin heavy and light chain expression on B cell differentiation. *Genes & Development*. 1994;**8**:1043-1057
- [66] Melchers F et al. Roles of IgH and L chains and of surrogate H and L chains in the development of cells of the B lymphocyte lineage. *Annual Review of Immunology*. 1994;**12**:209-225
- [67] Herzog S, Reth M, Jumaa H. Regulation of B cell proliferation and differentiation by pre-B-cell receptor signalling. *Nature Reviews. Immunology*. 2009;**9**:195-205
- [68] Ehlich A, Schaal S, Gu H, et al. Immunoglobulin heavy and light chain genes rearrange independently at early stages of B cell development. *Cell*. 1993;**72**:695-704
- [69] Li YS, Hayakawa K, Hardy RR. The regulated expression of B lineage associated genes during B cell differentiation in bone marrow and fetal liver. *The Journal of Experimental Medicine*. 1993;**178**:951-960
- [70] Shapiro AM, Schlissel MS, Baltimore D, Defranco AL. Stimulation of K light-chain gene rearrangement by the immunoglobulin  $\mu$  heavy chain in a pre-B-cell line. *Molecular and Cellular Biology*. 1993;**13**:5679-5690
- [71] Maki R, Kearney J, Paige C, Tonegawa S. Immunoglobulin gene rearrangement in immature B cells. *Science*. 1980;**209**:1366-1369
- [72] Perry RP, Kelley DE, Coleclough C, Kearney JF. Organization and expression of immunoglobulin genes in fetal liver hybridomas. *Proceedings of the National Academy of Sciences of the United States of America*. 1981;**78**:247-251
- [73] Koralov SB, Novobrantseva TI, Königsmann J, et al. Antibody repertoires generated by VH replacement and direct VH to JH joining. *Immunity*. 2006;**25**:43-53

- [74] Lutz J, Müller W, Jäck HM. VH replacement rescues progenitor B cells with two non-productive VDJ alleles. *Journal of Immunology*. 2006;**177**:7007-7714
- [75] Kleinfeld R, Hardy RR, Tarlinton D, et al. Recombination between an expressed immunoglobulin heavy-chain gene and a germline variable gene segment in a Ly 1+ B-cell lymphoma. *Nature*. 1986;**322**:843-846
- [76] Reth M, Gehrmann P, Petrac E, Wiese P. A novel VH to VHDJH joining mechanism in heavy-chain-negative (null) pre-B cells results in heavy-chain production. *Nature*. 1986;**322**:840-842
- [77] Sun A, Novobrantseva TI, Coffrea M, et al. VH replacement in primary immunoglobulin repertoire diversification. *Proceedings of the National Academy of Sciences of the United States of America*. 2015;**112**:E458-E466
- [78] Jung D, Giallourakis C, Mostoslavsky R, Alt FW. Mechanism and control of V(D)J recombination at the immunoglobulin heavy chain locus. *Annual Review of Immunology*. 2006;**24**:541-570
- [79] Akira S, Okazaki K, Sakano H. Two pairs of recombination signals are sufficient to cause V-(D)-J joining. *Science*. 1987;**238**:1134-1138
- [80] Schatz DG, Oettinger MA, Baltimore D. The V(D)J recombination-activating gene, RAG-1. *Cell*. 1989;**59**:1035-1048
- [81] Oettinger MA, Schatz DG, Gorka C, Baltimore D. Rag-1 and Rag-2, adjacent genes that synergistically activate V(D)J recombination. *Science*. 1990;**248**:1517-1523
- [82] van Gent DC, Ramsden DA, Gellert M. The RAG1 and RAG2 proteins establish the 12/23 rule in V(D)J recombination. *Cell*. 1996;**85**:107-113
- [83] Sleckman BP, Gorman JR, Alt FW. Accessibility control of antigen-receptor variable-region gene assembly: Role of cis-acting elements. *Annual Review of Immunology*. 1996;**14**:459-481
- [84] Klein U, Tu Y, Stolovitzky GA, et al. Transcriptional analysis of the B cell germinal center reaction. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;**100**:2639-2644
- [85] Matthias P, Rolink AG. Transcriptional networks in developing and mature B cells. *Nature Reviews. Immunology*. 2005;**5**:497-508
- [86] Shaffer AL, Rosenwald A, Hurt EM, et al. Signatures of the immune response. *Immunity*. 2001;**15**:375-385
- [87] Heinz S, Benner C, Spann N, et al. Simple combinations of lineage determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Molecular Cell*. 2010;**38**:576-589
- [88] Mercer EM, Lin YC, Benner C, et al. Multilineage priming of enhancer repertoires precedes commitment to the B and myeloid cell lineages in hematopoietic progenitors. *Immunity*. 2011;**35**:413-425

- [89] Iwafuchi-Doi M, Zaret KS. Pioneer transcription factors in cell reprogramming. *Genes & Development*. 2014;**28**:2679-2692
- [90] Rothenberg EV. Transcriptional control of early T and B cell developmental choices. *Annual Review of Immunology*. 2014;**32**:283-321
- [91] Boller S, Grosschedl R. The regulatory network of B-cell differentiation: A focused view of early B-cell factor 1 function. *Immunological Reviews*. 2014;**261**:102-115
- [92] Nutt SL, Kee BL. The transcriptional regulation of B cell lineage commitment. *Immunity*. 2007;**26**:715-725
- [93] Costa G, Kouskoff V, Lacaud G. Origin of blood cells and HSC production in the embryo. *Trends in Immunology*. 2012;**35**:215-223
- [94] Kim I, Saunders TL, Morrison SJ. Sox17 dependence distinguishes the transcriptional regulation of fetal from adult hematopoietic stem cells. *Cell*. 2007;**130**:470-483
- [95] Cuvertino S, Lacaud G, Kouskoff V. SOX7-enforced expression promotes the expansion of adult blood progenitors and blocks B-cell development. *Open Biology*. 2016;**6**:1-12
- [96] Stehling-Sun S, Dade J, Nutt SL, DeKoter RP, Camargo FD. Regulation of lymphoid versus myeloid fate 'choice' by the transcription factor Mef2c. *Nature Immunology*. 2009;**10**:289-296
- [97] Wang W, Org T, Montel-Hagen A, et al. MEF2C protects bone marrow B-lymphoid progenitors during stress haematopoiesis. *Nature Communications*. 2016;**7**:1-15
- [98] Kurkewich JL, Klopfenstein N, Hallas WM, et al. Arid3b is critical for B lymphocyte development. *PLoS One*. 2016;**11**:1-19
- [99] de Pooter RF, Kee BL. E proteins and the regulation of early lymphocyte development. *Immunological Reviews*. 2010;**238**:93-109
- [100] Dias S, Mansson R, Gurbuxani S, Sigvardsson M, Kee BL. E2A proteins promote development of lymphoid-primed multipotent progenitors. *Immunity*. 2008;**29**:217-227
- [101] Bain G, Maandag EC, Izon DJ, et al. E2A proteins are required for proper B cell development and initiation of immunoglobulin gene rearrangements. *Cell*. 1994;**79**:885-892
- [102] Schwickert TA, Tagoh H, Gultekin S, et al. Stage-specific control of early B cell development by the transcription factor Ikaros. *Nature Immunology*. 2014;**15**:283-293
- [103] Wang JH, Nichogiannopoulou A, Wu L, et al. Selective defects in the development of the fetal and adult lymphoid system in mice with an Ikaros null mutation. *Immunity*. 1996;**5**:537-549
- [104] Dakic A, Metcalf D, Di Rago L, et al. PU.1 regulates the commitment of adult hematopoietic progenitors and restricts granulopoiesis. *The Journal of Experimental Medicine*. 2005;**201**:1487-1502



- [105] Scott EW, Simon MC, Anastasi J, Singh H. Requirement of transcription factor PU.1 in the development of multiple hematopoietic lineages. *Science*. 1994;**265**:1573-1577
- [106] DeKoter RP, Singh H. Regulation of B lymphocyte and macrophage development by graded expression of PU.1. *Science*. 2000;**288**:1439-1441
- [107] Spooner CJ, Cheng JX, Pujadas E, Laslo P, Singh H. A recurrent network involving the transcription factors PU.1 and Gfi1 orchestrates innate and adaptive immune cell fates. *Immunity*. 2009;**31**:576-586
- [108] Zou GM, Chen JJ, Yoder MC, Wu W, Rowley JD. Knockdown of Pu.1 by small interfering RNA in CD34+embryoid body cells derived from mouse ES cells turns cell fate determination to pro-B cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;**102**:13236-13241
- [109] Kee BL, Murre C. Induction of early B cell factor (EBF) and multiple B lineage genes by the basic helix-loop-helix transcription factor E12. *The Journal of Experimental Medicine*. 1998;**188**:699-713
- [110] Lin H, Grosschedl R. Failure of B-cell differentiation in mice lacking the transcription factor EBF. *Nature*. 1995;**376**:263-267
- [111] Lin YC, Jhunjhunwala S, Benner C, et al. A global network of transcription factors, involving E2A, EBF1 and Foxo1, that orchestrates B cell fate. *Nature Immunology*. 2010;**11**:635-643
- [112] Singh H, Medina KL, Pongubala JM. Contingent gene regulatory networks and B cell fate specification. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;**102**:4949-4953
- [113] Batliner J, Mancarelli MM, Jenal M, et al. CLEC5A (MDL-1) is a novel PU.1 transcriptional target during myeloid differentiation. *Molecular Immunology*. 2011;**48**:714-719
- [114] Fraszczak J, Helness A, Chen R, et al. Threshold levels of Gfi1 maintain E2A activity for B cell commitment via repression of Id1. *PLoS One*. 2016;**11**:e0160344
- [115] Milanovic M, Heise N, De Silva NS, et al. Differential requirements for the canonical NF- $\kappa$ B transcription factors c-REL and RELA during the generation and activation of mature B-cells. *Immunology and Cell Biology*. 2017;**95**:261-271
- [116] Shinohara H, Nagashima T, Cascalho MI, Kurosaki T. TAK1 maintains the survival of immunoglobulin k-chain-positive B cells. *Genes to Cells*. 2016;**21**:1233-1243
- [117] Derudder E, Herzog S, Labic V, et al. Canonical NF- $\kappa$ B signaling is uniquely required for the long-term persistence of functional mature B cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2016;**113**:5065-5070
- [118] De Silva NS, Anderson MM, Carette A, et al. Transcription factors of the alternative NF- $\kappa$ B pathway are required for germinal center B-cell development. *Proceedings of the National Academy of Sciences of the United States of America*. 2016;**113**:9063-9068

- [119] Lu R, Medina KL, Lancki DW, Singh H. IRF-4,8 orchestrate the pre-B-to-B transition in lymphocyte development. *Genes & Development*. 2003;**17**:1703-1708
- [120] Piovesan D, Tempany J, Di Pietro A, et al. C-Myb regulates the T-bet-dependent differentiation program in B cells to coordinate antibody responses. *Cell Reports*. 2017;**19**:461-470
- [121] Schlissel M, Voronova A, Baltimore D. Helix-loop-helix transcription factor E47 activates germ-line immunoglobulin heavy-chain gene transcription and rearrangement in a pre-T-cell line. *Genes & Development*. 1991;**5**:1367-1376
- [122] Vu TT, Gatto D, Turner V, et al. Impaired B cell development in the absence of Krüppel-like factor 3. *Journal of Immunology*. 2011;**187**:5032-5042
- [123] Zhang XK, Moussa O, LaRue A, et al. The transcription factor Fli-1 modulates marginal zone and follicular B cell development in mice. *Journal of Immunology*. 2008;**181**:1644-1654
- [124] Orlanskia S, Labib V, Reizela Y, et al. Tissue-specific DNA demethylation is required for proper B-cell differentiation and function. *Proceedings of the National Academy of Sciences of the United States of America*. 2016;**113**:5018-5023
- [125] Zhangl S, Zhu I, Deng T, et al. HMGN proteins modulate chromatin regulatory sites and gene expression during activation of naïve B cells. *Nucleic Acids Research*. 2016;**44**:7144-7158
- [126] Boller S, Ramamoorthy S, Akbas D, et al. Pioneering activity of the C-terminal domain of EBF1 shapes the chromatin landscape for B cell programming. *Immunity*. 2016;**44**:527-541
- [127] Pongubala JM, Northrup DL, Lancki DW, et al. Transcription factor EBF restricts alternative lineage options and promotes B cell fate commitment independently of Pax5. *Nature Immunology*. 2008;**9**:203-215
- [128] Zandi S, Mansson R, Tsapogas P, et al. EBF1 is essential for B-lineage priming and establishment of a transcription factor network in common lymphoid progenitors. *Journal of Immunology*. 2008;**181**:3364-3372
- [129] Treiber T, Mandel EM, Pott S, et al. Early B cell factor 1 regulates B cell gene networks by activation, repression, and transcription-independent poising of chromatin. *Immunity*. 2010;**32**:714-725
- [130] Banerjee A, Northrup D, Boukarabila H, Jacobsen SE, Allman D. Transcriptional repression of Gata3 is essential for early B cell commitment. *Immunity*. 2013;**38**:930-942

